

EXHIBIT 22

(12) **United States Patent**
Fodor et al.(10) **Patent No.:** **US 6,440,667 B1**
(45) **Date of Patent:** **Aug. 27, 2002**(54) **ANALYSIS OF TARGET MOLECULES USING AN ENCODING SYSTEM**(75) Inventors: **Stephen P. A. Fodor**, Palo Alto;
Dennis W. Solas, San Francisco;
William J. Dower, Menlo Park, all of
CA (US)(73) Assignee: **Affymetrix Inc.**, Santa Clara, CA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.(21) Appl. No.: **09/362,089**(22) Filed: **Jul. 28, 1999****Related U.S. Application Data**(60) Division of application No. 09/056,927, filed on Apr. 8,
1998, now Pat. No. 6,197,506, which is a continuation of
application No. 08/670,118, filed on Jun. 25, 1996, now Pat.
No. 5,800,992, which is a division of application No.
08/168,904, filed on Dec. 15, 1993, now abandoned, which
is a continuation of application No. 07/624,114, filed on
Dec. 6, 1990, now abandoned, which is a continuation-in-
part of application No. 07/492,462, filed on Mar. 7, 1990,
now Pat. No. 5,143,854, and a continuation-in-part of appli-
cation No. 07/362,901, filed on Jun. 7, 1989, now aban-
doned.(51) **Int. Cl.**⁷ **C12Q 1/68**; G01N 33/53;
C12M 1/34; C07H 21/04; A61K 38/00(52) **U.S. Cl.** **435/6**; 435/7.1; 435/287.2;
435/288.3; 530/300; 530/350; 530/387.1;
536/23.1; 536/24.3(58) **Field of Search** 435/6, 7.1, 287.2,
435/288.3; 536/23.1, 24.3; 530/350, 300,
387.1(56) **References Cited****U.S. PATENT DOCUMENTS**3,730,844 A 5/1973 Gilham et al. 195/103.5 R
3,849,137 A 11/1974 Barzynski et al. 96/97
3,862,056 A 1/1975 Hartman 252/511

(List continued on next page.)

FOREIGN PATENT DOCUMENTSCA 1284931 6/1991
DE 2242394 3/1974
DE 3440141 5/1986

(List continued on next page.)

OTHER PUBLICATIONSBrenner et al. "In vitro cloning of complex mixtures of DNA
on microbeads: Physical separation of differentially
expressed cDNAs" Proc Natl Acad Sci USA 2000,
97:1665-1670.Brenner et al. "Gene expression analysis by massively
parallel signature sequencing (MPSS) on microbead arrays"
Nature Biotechnology 2000, 18:630-634.Tyagi "Taking a census of mRNA populations with micro-
beads" Nature Biotechnology 2000, 18:597-598.3,939,350 A 2/1976 Kronick et al. 250/365
4,072,576 A 2/1978 Arwin et al. 195/103.5 R
4,121,222 A 10/1978 Diebold et al. 347/7
4,180,739 A 12/1979 Abu-Shumays 250/461 R
4,216,245 A 8/1980 Johnson 427/2.13
4,238,757 A 12/1980 Schenck 357/25
4,269,933 A 5/1981 Pazos 430/291
4,314,821 A 2/1982 Rice 23/230 B
4,327,073 A 4/1982 Huang 424/1
4,339,528 A 7/1982 Goldman 430/323
4,342,905 A 8/1982 Fujii et al. 250/201
4,373,071 A 2/1983 Itakura 525/375
4,395,486 A 7/1983 Wilson et al. 435/6
4,405,771 A 9/1983 Jagur 528/266
4,444,878 A 4/1984 Paulus 435/7
4,444,892 A 4/1984 Malmros 436/528
4,448,534 A 5/1984 Wertz et al. 356/435
4,458,066 A 7/1984 Caruthers et al. 536/27
4,483,920 A 11/1984 Gillespie et al. 435/6
4,500,707 A 2/1985 Caruthers et al. 536/27
4,500,919 A 2/1985 Schreiber 358/78
4,516,833 A 5/1985 Fusek 350/162.12
4,517,338 A 5/1985 Urdea et al. 525/54.11
4,533,682 A 8/1985 Tortorello et al. 523/414
4,537,861 A 8/1985 Elings et al. 436/518
4,542,102 A 9/1985 Dattagupta et al. 435/6
4,555,490 A 11/1985 Merrill 436/86
4,556,643 A 12/1985 Paau et al. 435/5
4,562,157 A 12/1985 Lowe et al. 435/291
4,563,419 A 1/1986 Ranki et al. 435/6
4,569,967 A 2/1986 Kornreich et al. 525/54.11
4,580,895 A 4/1986 Patel 356/39
4,584,277 A 4/1986 Ullman 436/501
4,588,682 A 5/1986 Groet et al. 435/6
4,591,570 A 5/1986 Chang 435/7.24
4,598,049 A 7/1986 Zelinka et al. 422/116
4,613,566 A 9/1986 Potter 435/6
4,624,915 A 11/1986 Schindler et al. 435/4
4,626,684 A 12/1986 Landa 250/328
4,631,211 A 12/1986 Houghten 428/35
4,637,861 A 1/1987 Krull et al. 204/1 T
4,656,127 A 4/1987 Mundy 435/6
4,670,380 A 6/1987 Dattagupta 435/6
4,677,054 A 6/1987 White et al. 435/6
4,681,859 A 7/1987 Kramer 436/501
4,683,195 A 7/1987 Mullis et al. 435/6
4,683,202 A 7/1987 Mullis 435/91
4,689,405 A 8/1987 Frank et al. 536/27
4,704,353 A 11/1987 Humphries et al. 435/4
4,711,955 A 12/1987 Ward et al. 536/29
4,713,326 A 12/1987 Dattagupta et al. 435/6
4,713,347 A 12/1987 Mitchell et al. 436/501
4,715,413 A 12/1987 Backlund et al. 141/94

(List continued on next page.)

Primary Examiner—Stephanie Zitomer(74) *Attorney, Agent, or Firm*—Morgan, Lewis & Bockius
LLP(57) **ABSTRACT**The present invention provides methods and apparatus for
sequencing, fingerprinting and mapping biological
macromolecules, typically biological polymers. The meth-
ods make use of a plurality of sequence specific recognition
reagents which can also be used for classification of bio-
logical samples, and to characterize their sources.**53 Claims, 2 Drawing Sheets**

US 6,440,667 B1

Page 2

U.S. PATENT DOCUMENTS

4,716,106 A	12/1987	Chiswell	435/6	5,096,807 A	3/1992	Leaback	435/6
4,719,179 A	1/1988	Barany	435/172.1	5,100,626 A	3/1992	Levin	422/100
4,719,615 A	1/1988	Feyrer et al.	369/284	5,100,777 A	3/1992	Chang	435/7.24
4,722,906 A	2/1988	Guire	436/501	5,112,962 A	5/1992	Letsinger et al.	536/27
4,728,502 A	3/1988	Hamill	422/116	5,141,813 A	8/1992	Nelson	428/402
4,728,591 A	3/1988	Clark et al.	430/5	5,143,854 A	9/1992	Pirrung et al.	436/518
4,731,325 A	3/1988	Palva et al.	435/6	5,149,625 A	9/1992	Church et al.	435/6
4,737,344 A	4/1988	Koizumi et al.	422/100	5,153,319 A	10/1992	Caruthers et al.	536/27
4,755,458 A	7/1988	Rabbani et al.	435/5	5,164,319 A	11/1992	Hafeman et al.	435/287.1
4,762,881 A	8/1988	Kauer	525/54.11	5,171,695 A	12/1992	Ekins	436/501
4,766,062 A	8/1988	Diamond et al.	435/6	5,188,963 A	2/1993	Stapleton	435/288.3
4,767,700 A	8/1988	Wallace	435/6	5,192,980 A	3/1993	Dixon et al.	356/326
4,777,019 A	10/1988	Dandekar	422/68	5,200,051 A	4/1993	Cozzette et al.	204/403
4,780,504 A	10/1988	Buendia et al.	525/54.11	5,202,231 A	4/1993	Drmanac et al.	435/6
4,786,170 A	11/1988	Groeblor	356/318	5,206,137 A	4/1993	Ip et al.	435/6
4,786,684 A	11/1988	Glass	525/54.1	5,215,882 A	6/1993	Bahl et al.	435/6
4,794,150 A	12/1988	Steel	525/54.11	5,215,889 A	6/1993	Schultz	435/41
4,808,508 A	2/1989	Platzer	430/143	5,219,726 A	6/1993	Evans	435/6
4,810,869 A	3/1989	Yabe et al.	250/201	5,225,326 A	7/1993	Bresser et al.	435/6
4,811,062 A	3/1989	Tabata et al.	356/152	5,232,829 A	8/1993	Longiaru et al.	435/6
4,811,218 A	3/1989	Hunkapiller et al.	204/461	5,235,028 A	8/1993	Barany et al.	528/335
4,812,512 A	3/1989	Buendia et al.	525/54.11	5,242,974 A	9/1993	Holmes	525/54.11
4,820,630 A	4/1989	Taub	435/5	5,252,743 A	10/1993	Barrett et al.	548/303.7
4,822,566 A	4/1989	Newman	422/68	5,256,549 A	10/1993	Urdea et al.	435/91
4,833,092 A	5/1989	Geysen	436/501	5,258,506 A	11/1993	Urdea et al.	536/23.1
4,844,617 A	7/1989	Kelderman et al.	356/372	5,306,641 A	4/1994	Saccocio	436/85
4,846,552 A	7/1989	Veldkamp et al.	350/162.2	5,310,893 A	5/1994	Erlich et al.	536/24.31
4,849,513 A	7/1989	Smith et al.	536/27	5,324,633 A	6/1994	Fodor et al.	435/6
4,855,225 A	8/1989	Fung et al.	435/6	5,328,824 A	7/1994	Ward et al.	435/6
4,865,990 A	9/1989	Stead et al.	435/803	5,348,855 A	9/1994	Dattagupta	435/6
4,868,103 A	9/1989	Stavrianopoulos et al.	435/5	5,384,261 A	1/1995	Winkler et al.	436/518
4,874,500 A	10/1989	Madou et al.	204/412	5,405,783 A	4/1995	Pirrung et al.	436/518
4,877,745 A	10/1989	Hayes et al.	436/166	5,424,186 A	6/1995	Fodor et al.	435/6
4,886,741 A	12/1989	Schwartz	435/5	5,424,188 A	6/1995	Schneider et al.	435/6
4,888,278 A	12/1989	Singer et al.	435/6	5,432,099 A	7/1995	Ekins	436/518
4,921,805 A	5/1990	Gebeyehu et al.	435/270	5,436,327 A	7/1995	Southern et al.	536/25.34
4,923,901 A	5/1990	Koester et al.	521/53	5,445,934 A	8/1995	Fodor et al.	435/6
4,925,785 A	5/1990	Wang et al.	435/6	5,447,841 A	9/1995	Gray et al.	435/6
4,931,384 A	6/1990	Layton et al.	435/7.31	5,451,505 A	9/1995	Dollinger	
4,946,942 A	8/1990	Fuller et al.	530/335	5,474,796 A	12/1995	Brennan	427/2.13
4,965,188 A	10/1990	Mullis et al.	435/6	5,486,452 A	1/1996	Gordon et al.	435/5
4,973,493 A	11/1990	Guire	427/2	5,489,507 A	2/1996	Chehab	435/6
4,979,959 A	12/1990	Guire	623/66	5,489,678 A	2/1996	Fodor et al.	536/22.1
4,981,783 A	1/1991	Augenlicht	435/6	5,492,806 A	2/1996	Drmanac et al.	435/5
4,981,985 A	1/1991	Kaplan et al.	556/50	5,494,810 A	2/1996	Barany et al.	435/91.52
4,984,100 A	1/1991	Takayama et al.	360/49	5,510,270 A	4/1996	Fodor et al.	436/518
4,987,065 A	1/1991	Stavrianopoulos et al.	435/5	5,525,464 A	6/1996	Drmanac et al.	435/6
4,988,617 A	1/1991	Landegren et al.	435/6	5,527,681 A	6/1996	Holmes	435/6
4,992,383 A	2/1991	Farnsworth	436/89	5,552,270 A	9/1996	Khrapko et al.	435/6
4,994,373 A	2/1991	Stavrianopoulos et al.	435/6	5,556,961 A	9/1996	Foot et al.	536/27.1
5,002,867 A	3/1991	Macevitz	435/6	5,561,071 A	10/1996	Hollenberg et al.	437/1
5,006,464 A	4/1991	Chu et al.	435/7.1	5,565,324 A	10/1996	Still	
5,011,770 A	4/1991	Kung et al.	435/6	5,567,809 A	10/1996	Apple	
5,013,669 A	5/1991	Peters, Jr. et al.	436/518	5,569,584 A	10/1996	Augenlicht	435/6
5,021,550 A	6/1991	Zeiger	530/334	5,571,639 A	11/1996	Hubbell et al.	430/5
5,026,773 A	6/1991	Steel	525/54.11	5,573,905 A	11/1996	Lerner	
5,026,840 A	6/1991	Dattagupta et al.	536/27	5,593,839 A	1/1997	Hubbell et al.	435/6
5,028,525 A	7/1991	Gray et al.	435/6	5,599,720 A	2/1997	Ekins	436/501
5,028,545 A	7/1991	Soini	436/501	5,604,097 A	2/1997	Brenner	
5,037,882 A	8/1991	Steel	525/54.11	5,604,099 A	2/1997	Erlich et al.	435/6
5,043,265 A	8/1991	Tanke et al.	435/6	5,635,400 A	6/1997	Brenner	
5,047,524 A	9/1991	Andrus et al.	536/27	5,641,634 A *	6/1997	Mandecki	435/6
5,064,754 A	11/1991	Mills	435/6	5,643,728 A	7/1997	Slater et al.	435/6
5,075,077 A *	12/1991	Durley, III et al.	422/56	5,653,939 A	8/1997	Hollis et al.	422/50
5,077,085 A	12/1991	Schnur et al.	357/4	5,654,413 A	8/1997	Brenner	
5,077,210 A	12/1991	Eigler et al.	435/176	5,667,667 A	9/1997	Southern	205/687
5,079,600 A	1/1992	Schnur et al.	357/4	5,667,972 A	9/1997	Drmanac et al.	435/6
5,081,584 A	1/1992	Omichinski et al.	364/497	5,690,894 A	11/1997	Pinkel	
5,082,830 A	1/1992	Brakel et al.	514/44	5,695,940 A	12/1997	Drmanac et al.	435/6
5,091,652 A	2/1992	Mathies et al.	250/458.1	5,698,393 A	12/1997	Macioszek et al.	435/5
				5,700,637 A	12/1997	Southern	435/6

US 6,440,667 B1

Page 3

5,707,806	A	1/1998	Shuber	435/6	GB	2233654	1/1991
5,744,305	A	4/1998	Fodor et al.	435/6	GB	2248840	9/1992
5,751,629	A *	5/1998	Nova et al.	365/151	JP	49-110601	10/1974
5,770,367	A	6/1998	Southern		JP	60-248669	12/1985
5,776,737	A	7/1998	Dunn	435/91.1	JP	63-084499	4/1988
5,777,888	A	7/1998	Rine et al.	364/496	JP	63-223557	9/1988
5,800,992	A	9/1998	Fodor et al.	435/6	JP	1-233447	9/1989
5,804,563	A	9/1998	Still		NO	P 913186	8/1991
5,807,522	A	9/1998	Brown et al.	422/50	WO	WO 84/03151	8/1984
5,807,683	A	9/1998	Brenner		WO	WO 84/03564	9/1984
5,830,645	A	11/1998	Pinkel et al.	435/6	WO	WO 85/01051	3/1985
5,843,767	A	12/1998	Beattie	435/287.1	WO	WO 86/00991	2/1986
5,846,708	A	12/1998	Hollis et al.	435/6	WO	WO 86/06487	11/1986
5,846,719	A	12/1998	Brenner		WO	WO 88/01058	2/1988
5,863,722	A	1/1999	Brenner		WO	WO 88/04777	6/1988
5,869,237	A	2/1999	Ward et al.	435/6	WO	WO 89/05616	6/1989
5,871,697	A	2/1999	Rothberg et al.	422/68.1	WO	WO 89/08834	9/1989
5,972,619	A	10/1999	Drmanac et al.	435/6	WO	WO 89/10977	11/1989
6,018,041	A	1/2000	Drmanac et al.	536/24.3	WO	WO 89/11548	11/1989
6,023,540	A	2/2000	Walt		WO	WO 89/12819	12/1989
6,025,136	A	2/2000	Drmanac et al.	435/6	WO	WO 90/00626	1/1990
6,040,166	A	3/2000	Erllich et al.	435/194	WO	WO 90/00887	2/1990
6,054,270	A	4/2000	Southern		WO	WO 90/15070	2/1990
6,060,240	A	5/2000	Kamb		WO	WO 90/03382	4/1990

FOREIGN PATENT DOCUMENTS

DE	3505287	3/1988	WO	WO 90/04652	5/1990
DE	3722958	1/1989	WO	WO 90/05789	5/1990
EP	046 083	2/1982	WO	WO 90/07582	7/1990
EP	088 636	9/1983	WO	WO 91/00868	1/1991
EP	103 197	3/1984	WO	WO 91/04266	4/1991
EP	127 438	12/1984	WO	WO 91/07087	5/1991
EP	063 810	3/1986	WO	WO 92/16655	1/1992
EP	174 879	3/1986	WO	WO 92/10092	6/1992
EP	194 132	9/1986	WO	WO 92/10588	6/1992
EP	228 075	7/1987	WO	WO 93/02992	2/1993
EP	245 662	11/1987	WO	WO 93/09668	5/1993
EP	268 237	5/1988	WO	WO 88/01302	6/1993
EP	130 523	6/1988	WO	WO 93/11262	6/1993
EP	281 927	9/1988	WO	WO 93/17126	9/1993
EP	228 310	10/1988	WO	WO 93/22456	11/1993
EP	288 310	10/1988	WO	WO 93/22480	11/1993
EP	304 202	2/1989	WO	WO 95/00530	1/1995
EP	307 476	3/1989	WO	WO 95/11995	5/1995
EP	319 012	6/1989	WO	WO 95/33846	12/1995
EP	328 256	8/1989	WO	WO 96/23078	8/1996
EP	333 561	9/1989	WO	WO 97/10365	3/1997
EP	337 498	10/1989	WO	WO 97/17317	5/1997
EP	386 229	4/1990	WO	WO 97/19410	5/1997
EP	373 203	6/1990	WO	WO 97/27317	7/1997
EP	392 546	10/1990	WO	WO 97/29212	8/1997
EP	142 299	12/1990	WO	WO 97/31256	8/1997
EP	173 339	1/1992	WO	WO 97/45559	12/1997
EP	171 150	3/1992	WO	WO 98/03673	1/1998
EP	237 362	3/1992	WO	WO 98/31836	7/1998
EP	185 547	6/1992	WO	WO 99/60007	11/1999
EP	260 634	6/1992	YU	P-570/87	4/1987
EP	232 967	4/1993	YU	18617/87	9/1987
EP	235 726	5/1993			
EP	476 014	8/1994			
EP	225 807	10/1994			
EP	717 113	6/1996			
EP	721 016	7/1996			
EP	535 242	9/1997			
EP	848 067	6/1998			
EP	619 321	1/1999			
FR	2559783	3/1988			
GB	2 129 551 *	5/1984			
GB	2156074	3/1988			
GB	2196476	4/1988			
GB	8810400.5	5/1988			

OTHER PUBLICATIONS

Miller et al. "Detection of bacteria by hybridization of rRNA with DNA-latex and immunodetection of hybrids" J Clin Microbiol 1988, 26:1271-1276.

Sequencing by Hybridization Workshop, listing of participants and workshop presentation summaries, from workshop held Nov. 19-20, 1991.

"A Sequencing Reality Check," *Science*, 242:1245 (1988).

"Affymax raises \$25 million to develop high-speed drug discovery system," *Biotechnology News*, 10(3):7-8 (1990).

"Preparation of fluorescent-labeled DNA and its use as a probe in molecular hybridization," *Bioorg Khim*, 12(11):1508-1513 (1986).

US 6,440,667 B1

Page 4

- Abbott et al., "Manipulation of the Wettability of Surfaces on the 0.1- to 1-Micrometer Scale Through Micromachining and Molecular Self-Assembly," *Science*, 257:1380-1382 (1992).
- Adams et al., "Complementary DNA Sequencing: Expressed Sequence Tags and Human Genome Project," *Science*, 252(5013):1651-1656 (1991).
- Adams et al., "Photolabile Chelators That "Cage" Calcium with Improved Speed of Release and Pre-Photolysis Affinity," *J. Gen. Physiol.*, p. 9a (12/86).
- Adams et al., "Biologically Useful Chelators That Take Up Ca²⁺ upon Illumination," *J. Am. Chem. Soc.*, 111:7957-7968 (1989).
- Ajayaghosh et al., "Solid-Phase Synthesis of N-Methyl- and N-Ethylamides of Peptides Using Photolytically Detachable ((3-Nitro-4(alkylamino)methyl)benzamido)methylpolystyrene Resin," *J. Org. Chem.*, 55(9):2826-2829 (1990).
- Ajayaghosh et al., "Solid-phase synthesis of C-terminal peptide amides using a photoremovable α -methylphenacylamido anchoring linkage," *Proc. Ind. Natl. Sci (Chem.Sci.)*, 100(5):389-396 (1988).
- Ajayaghosh et al., "Polymer-supported Solid-phase Synthesis of C-Terminal Peptide N-Methylamides Using a Modified Photoremovable 3-Nitro-4-N-methylaminomethylpolystyrene Support," *Ind.J.Chem.*, 27B:1004-1008 (1988).
- Ajayaghosh et al., "Polymer-Supported Synthesis of Protected Peptide Segments on a Photosensitive o-Nitro (α -Methyl)Bromobenzyl Resin," *Tetrahedron*, 44(21):6661-6666 (1988).
- Amit et al., "Photosensitive Protecting Groups of Amino Sugars and Their Use in Glycoside Synthesis. 2-Nitrobenzyloxycarbonylamino and 6-Nitroveratryloxycarbonylamino Derivatives," *J.Org.Chem*, 39(2):192-196 (1974).
- Amit et al., "Photosensitive Protecting Groups—A Review," *Israel J. Chem.*, 12(1-2):103-113 (1974).
- Anand et al., "A 3.5 genome equivalent multi access YAC library: construction, characterisation, screening and storage," *Nuc. Acids Res.*, 18(8):1951-1956 (1990).
- Anderson et al., "Quantitative Filter Hybridisation," chapter 3 from *Nucleic Acid Hybridization a practical approach*, pp. 73-111, Hames et al., eds., IRL Press (1985).
- Applied Biosystems, Model 431A Peptide Synthesizer User's manual, Sections 2 and 6, (Aug. 15, 1989).
- Arnold et al., "A Novel Universal Support for DNA & RNA Synthesis," abstract from *Federation Proceedings*, 43(7): abstract No. 3669 (1984).
- Atherton et al., *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, (1989) tbl. of cont., pp. vii-ix.
- Augenlicht et al., "Cloning and Screening of Sequences Expressed in a Mouse Colon Tumor," *Cancer Research*, 42:1088-1093 (1982).
- Augenlicht et al., "Expression of Cloned Sequences in Biopsies of Human Colonic Tissue and in Colonic Carcinoma Cells Induced to Differentiate in Vitro," *Cancer Res.*, 47:6017-6021 (1987).
- Bains, W., "Hybridization Methods for DNA Sequencing," *Genomics*, 11(2):294-301 (1991).
- Bains et al., "A Novel Method for Nucleic Acid Sequence Determination," *J.Theor.Biol.*, 135:303-307 (1988).
- Bains, W., "Alternative Routes Through the Genome," *Biotechnology*, 8:1251-1256 (1988).
- Balachander et al., "Functionalized Siloxy-Anchored Monolayers with Exposed Amino, Azido, Bromo, or Cyano Groups," *Tetrahed. Ltrs.*, 29(44):5593-5594 (1988).
- Baldwin et al., "New Photolabile Phosphate Protecting Groups," *Tetrahed.*, 46(19):6879-6884 (1990).
- Bannwarth et al., "Laboratory Methods, A System for the Simultaneous Chemical synthesis of Different DNA Fragments on Solid Support," *DNA*, 5(5):413-419 (1986).
- Bannwarth, W., "Gene Technology: a Challenge for a Chemist," *CHIMIA*, 41(9):302-317 (1987).
- Barany, F., "Genetic disease detection and DNA amplification using cloned thermostable ligase," *PNAS*, 88:189-193 (1991).
- Barltrop et al., "Photosensitive Protective Groups," *Chemical Communications*, pp. 822-823 (1966).
- Barinaga, M., "Will 'DNA Chip' Speed Genome Initiative," *Science*, 253:1489 (1985).
- Bart et al., "Microfabricated Electrohydrodynamic Pumps," *Sensors and Actuators*, A21-A23:193-197 (1990).
- Bartsh et al., "Cloning of mRNA sequences from the human colon: Preliminary characterisation of defined mRNAs in normal and neoplastic tissues," *Br.J.Can.*, 54:791-798 (1986).
- Baum, R., "Fledgling firm targets drug discovery process," *Chem. Eng. News*, p. 10-11 (1990).
- Beltz et al., "Isolation of Multigene Families and Determination of Homologies by Filter Hybridization Methods," *Methods in Enzymology*, 100:266-285 (1983).
- Benschop, Chem. Abstracts 114(26):256643 (1991).
- Bhatia et al., "New Approach To Producing Patterned Biomolecular Assemblies," *J. American Chemical Society*, 114:4432-4433 (1992).
- Biorad Chromatography Electrophoresis Immunochemistry Molecular Biology HPLC catalog M 1987 pp. 182.
- Blawas et al., "Step-and-Repeat Photopatterning of Protein Features Using Caged-Biotin-BSA: Characterization and Resolution," *Langmuir*, 14(15):4243-4250 (1998).
- Blawas, A.S., "Photopatterning of Protein Features using Caged-biotin-Bovine Serum Albumin," dissertation for Ph.D at Duke University in 1998.
- Bos et al., "Amino-acid substitutions at codon 13 of the N-ras oncogene in human acute myeloid leukaemia," *Nature*, 315:726-730 (1985).
- Boyle et al., "Differential distribution of long and short interspersed element sequences in the mouse genome: Chromosome karyotyping by fluorescence in situ hybridization," *PNAS*, 87:7757-7761 (1990).
- Brock et al., "Rapid fluorescence detection of in situ hybridization with biotinylated bovine herpesvirus-1 DNA probes," *J.Veterinary Diagnostic Invest.*, 1:34-38 (1989).
- Burgi et al., "Optimization in Sample Stacking for High-Performance Capillary Electrophoresis," *Anal. Chem.*, 63:2042-2047 (1991).
- Cameron et al., "Photogeneration of Organic Bases from o-Nitrobenzyl-Derived Carbamates," *J. Am. Chem. Soc.*, 113:4303-4313 (1991).
- Carrano et al., "A High-Resolution, Fluorescence-Based, Semiautomated Method for DNA Fingerprinting," *Genomics*, 4:129-136 (1989).
- Caruthers, M.H., "Gene Synthesis Machines: DNA Chemistry and Its Uses," *Science*, 230:281-285 (1985).
- Chatterjee et al., "Inducible Alkylation of DNA Using an Oligonucleotide-Quinone Conjugate," *Am. J. Chem. Soc.* 112:6397-6399 (1990).

US 6,440,667 B1

Page 5

- Chee et al., "Accessing Genetic Information with High-Density DNA Arrays," *Science*, 274:610-614 (1996).
- Chehab et al., "Detection of sickle cell anaemia mutation by colour DNA amplification," *Lancet*, 335:15-17 (1990).
- Chehab et al., "Detection of specific DNA sequences by fluorescence amplification: a color complementation assay," *PNAS*, 86:9178-9182 (1989).
- Chetverin et al., "Oligonucleotide Arrays: New Concepts and Possibilities," *Biotechnology*, 12:1093-1099 (1994).
- Church et al., "Multiplex DNA sequencing," *Science*, 240:185-188 (1988).
- Church et al., "Genomic sequencing," *PNAS*, 81:1991-1995 (1984).
- Corbett et al., "Reaction of Nitroso Aromatics with Glyoxylic Acid. A New Path to Hydroxamic Acids," *J. Org. Chem.*, 45:2834-2839 (1980).
- Coulson et al., "Toward a physical map of the genome of the nematode *Caenorhabditis elegans*," *PNAS*, 83:7821-7825 (1986).
- Craig et al., "Ordering of cosmid clones covering the Herpes simplex virus type 1 (HSV-1) genome: a test case for fingerprinting by hybridization," *Nuc. Acid. Res.*, 18(9):2653-2660 (1990).
- Cummings et al., "Photoactivable Fluorophores. 1. Synthesis and Photoactivation of o-Nitrobenzyl-Quenched Fluorescent Carbamates," *Tetrahedron Letters*, 29(1):65-68 (1988).
- Dattagupta et al., "Rapid identification of Microorganisms by Nucleic Acid Hybridization after Labeling the Test Sample," *Anal. Biochem.*, 177:85-89 (1989).
- Dattagupta et al., "Nucleic Acid Hybridization: a Rapid Method for the Diagnosis of Infectious Diseases," *Perspectives in Antiinfective Therapy*, eds. Jackson et al., pp. 241-247 (1988).
- Dower et al., "The Search for Molecular Diversity (II): Recombinant and Synthetic Randomized Peptide Libraries," *Ann. Rep. Med. Chem.*, 26:271-280 (1991).
- Diggelmann, "Investigating the VLSIPS synthesis process," Sep. 9, 1994.
- Di Mauro et al., "DNA Technology in Chip Construction," *Adv. Mater.*, 5(5):384-386 (1993).
- Drmanac et al., "An Algorithm for the DNA Sequence Generation from k-Tuple Word Contents of the Minimal Number of Random Fragments," *J. Biomol. Struct. Dyn.*, 8(5):1085-1102 (1991).
- Drmanac et al., "Partial Sequencing by Oligo-Hybridization Concept and Applications in Genome Analysis," 1st Int. Conf. Electrophor., Supercomp., Hum. Genome pp. 60-74 (1990).
- Drmanac et al., "Sequencing by Oligonucleotide Hybridization: A Promising Framework in Decoding of the Genome Program?," 1st Int. Conf. Electrophor., Supercomp., Hum. Genome pp. 47-59 (1990).
- Drmanac et al., "Laboratory Methods, Reliable Hybridization of Oligonucleotides as Short as Six Nucleotides," *DNA and Cell Biol.*, 9(7):527-534 (1990).
- Drmanac et al., "Sequencing of Megabase Plus DNA by Hybridization: theory of the Method," *Genomics*, 4:114-128 (1989).
- Drmanac et al., "Sequencing of Megabase Plus DNA by Hybridization: Theory of the Method," abstract of presentation given at Cold Spring Harbor Symposium on Genome Mapping and Sequencing, Apr. 27, 1988 thru May 1, 1988.
- Dulcey et al., "Deep UV Photochemistry of Chemisorbed Monolayers: Patterned Coplanar Molecular Assemblies," *Science*, 252:551-554 (1991).
- Duncan et al., "Affinity Chromatography of a Sequence-Specific DNA Binding Protein Using Teflon-Linked Oligonucleotides," *Analytical Biochemistry*, 169:104-108 (1988).
- Effenhauser et al., "Glass Chips for High-speed Capillary Electrophoresis Separations with Submicrometer Plate Heights," *Anal. Chem.*, 65:2637-2642 (1993).
- Effenhauser et al., "High-Speed Separation of Antisense Oligonucleotides on a Micromachined Capillary Electrophoresis Device," *Anal. Chem.*, 66:2949-2953 (1994).
- Ekins et al., "High Specific Activity Chemiluminescent and Fluorescent Markers: their Potential Application to High Sensitivity and 'Multi-analyte' Immunoassays," *J. Bioluminescence Chemiluminescence*, 4:59-78 (1989).
- Ekins et al., "Development of Microspot Multi-Analyte Ratiometric Immunoassay Using dual Fluorescent-Labelled Antibodies," *Anal. Chemica Acta*, 227:73-96 (1989).
- Ekins et al., "Multianalyte Microspot Immunoassay-Microanalytical 'Compact Disk' of the Future," *Clin. Chem.*, 37(11):1955-1967 (1991).
- Ekins, R.P., "Multi-Analyte immunoassay*," *J. Pharmaceutical Biomedical Analysis*, 7(2):155-168 (1989).
- Ekins et al., "Fluorescence Spectroscopy and its Application to a New Generation of High Sensitivity, Multi-Microspot, Multianalyte, Immunoassay," *Clin. Chim. Acta*, 194:91-114 (1990).
- Elder, J.K., "Analysis of DNA Oligonucleotide Hybridization Data by Maximum Entropy," in *Maximum Entropy and Bayesian Methods*, eds. Mohammad-Djafari and Dement, Kluwer, Dordrecht, pp. 363-371 (1992).
- Ellis, R.W., "The Applications of Synthetic Oligonucleotides to Molecular Biology," *Pharmaceutical Research*, 3(4):195-207 (1986).
- Evans et al., "Microfabrication for Automation of Molecular processes in Human Genome Analysis," *Clin. Chem.*, 41(11):1681 (1995).
- Evans et al., "Physical mapping of complex genomes by cosmid multiplex analysis," *PNAS*, 86:5030-5034 (1989).
- Ezaki et al., "Small-Scale DNA Preparation for Rapid Genetic Identification of *Campylobacter* Species without Radioisotope," *Microbiol. Immunology*, 32(2):141-150 (1988).
- Fan et al., "Mapping small DNA sequences by fluorescence in situ hybridization directly on banded metaphase chromosomes," *PNAS*, 87(16):6223-6227 (1990).
- Fan et al., "Micromachining of Capillary Electrophoresis Injectors and Separators on Glass Chips and Evaluation of Flow at Capillary Intersections," *Anal. Chem.*, 66:177-184 (1994).
- Feinberg et al., ADDENDUM to "A technique for Radiolabeling DNA Restriction Endonuclease Fragments to High Specific Activity," *Anal. Biochem.*, 137:266-267 (1984).
- Fettingner et al., "Stacked modules for micro flow systems in chemical analysis: concept and studies using an enlarged model," *Sensors and Actuators*, B17:19-25 (1993).
- Flanders et al., "A new interferometric alignment technique," *App. Phys. Ltrs.*, 31(7):426-429 (1977).
- Fodor et al., "Multiplexed biochemical assays with biological chips," *Nature*, 364:555-556 (1993).
- Fodor et al., "Light-directed, Spatially Addressable Parallel Chemical Synthesis," *Science*, 251:767-773 (1991).

US 6,440,667 B1

Page 6

- Forman et al., "Thermodynamics of Duplex Formation and Mismatch Discrimination on Photolithographically Synthesized Oligonucleotide Arrays," chapter 13pgs. 206-228 from *Molecular Modeling of Nucleic Acids*, ACS Symposium Series 682, Apr. 13-17, 1997, Leontis et al., eds.
- Frank et al., "Simultaneous Multiple Peptide Synthesis Under Continuous flow Conditions on Cellulose Paper Discs as Segmental Solid Supports," *Tetrahedron*, 44(19):6031-6040 (1988).
- Frank et al., "Automation of DNA Sequencing Reactions and Related Techniques: A Workstation for Micromanipulation of Liquids," *Bio/Technology*, 6:1211-1212 (1988).
- Frank et al., "Simultaneous Synthesis and Biological Applications of DNA Fragments: An Efficient and Complete Methodology," *Methods in Enzymology*, 154:221-250 (1987).
- Fuhr et al., "Travelling wave-driven microfabricated electrohydrodynamic pumps for liquids," *J. Micromech. Microeng.*, 4:217-226 (1994).
- Fuller et al., "Urethane-Protected Amino Acid N-Carboxy Anhydrides and Their Use in Peptide Synthesis," *J. Amer. Chem. Soc.*, 112(20):7414-7416 (1990).
- Furka et al., "General method for rapid synthesis of multi-component peptide mixtures," *Int. J. Peptide Protein Res.*, 37:487-493 (1991).
- Furka et al., "Cornucopia of Peptides by Synthesis," 14th Int. Congress of Biochem. abst.# FR:013, Jul. 10-15, 1988 Prague, Czechoslovakia.
- Furka et al., "More Peptides by Less Labour," abst. 288, Int. Symp. Med. Chem., Budapest Hungary Aug. 15-19, 1988. Gait, eds., pp. 1-115 from *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, (1984).
- Gazard et al., "Lithographic Technique Using Radiation-Induced Grafting of Acrylic Acid into Poly(Methyl Methacrylate) Films," *Polymer Engineering and Science*, 20(16):1069-1072 (1980).
- Gergen et al., "Filter replicas and permanent collections of recombinant DNA plasmids," *Nuc.Acids Res.*, 7(8):2115-2137 (1979).
- Getzoff et al., "Mechanisms of Antibody Binding to a Protein," *Science*, 235:1191-1196 (1987).
- Geysen et al., "Strategies for epitope analysis using peptide synthesis," *J. Immunol. Meth.*, 102:259-274 (1987).
- Geysen et al., "Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid," *PNAS*, 81:3998-4002 (1984).
- Geysen et al., "A synthetic strategy for epitope mapping," from *Peptides:Chem. & Biol.*, Proc. of 10th Am. Peptide Symp., May 23-28, 1987, pp. 519-523, (1987).
- Geysen, "Antigen-antibody interactions at the molecular level: adventures in peptide synthesis," *Immunol. Today*, 6(12):364-369 (1985).
- Geysen et al., "Cognitive Features of Continuous Antigenic Determinants," from *Synthetic Peptides: Approaches to Biological Probes*, pp. 19-30, (1989).
- Geysen et al., "Chemistry of Antibody Binding to a Protein," *Science*, 235:1184-1190 (1987).
- Geysen et al., "The delineation of peptides able to mimic assembled epitopes," 1986 CIBA Symp., pp. 130-149.
- Geysen et al., "Cognitive Features of Continuous Antigenic Determinants," *Mol. Recognit.*, 1(1):1-10 (1988).
- Geysen et al., "A Prio Ri Delineation of a Peptide Which Mimics A Discontinuous Antigenic Determinant," *Mol. Immunol.*, 23(7):709-715 (1986).
- Ghosh et al., "Covalent attachment of oligonucleotides to solid supports," *Nuc. Acids Res.*, 15(13):5353-5373 (1987).
- Gilon et al., "Backbone Cyclization: A New Method for Conferring Conformational Constraint on Peptides," *Biopolymers*, 31(6):745-750 (1991).
- Gingeras et al., "Hybridization properties of immobilized nucleic acids," *Nuc. Acids Res.*, 15(13):5373-5390 (87).
- Gummerlock et al., "RAS Enzyme-Linked Immunoblot Assay Discriminates p21 Species: A Technique to Dissect Gene Family Expression," *Anal. Biochem.*, 180:158-168 (1989).
- Gurney et al., "Activation of a potassium current by rapid photochemically generated step increases of intracellular calcium in rat sympathetic neurons," *PNAS*, 84:3496-3500 (1987).
- Haase et al., "Detection of Two Viral Genomes in Single Cells by Double-Label Hybridization in Situ and Color Microradioautography," *Science*, 227:189-192 (1985).
- Hacia, et al., "Two color hybridization analysis using high density oligonucleotide arrays and energy transfer dyes," *Nuc. Acids Res.*, 26(16):3865-3866 (1998).
- Hack, M.L., "Conics Formed to Make Fluid & Industrial Gas Micromachines," *Genetic Engineering News*, 15(18):1, 29 (1995).
- Hagedorn et al., "Pumping of Water Solutions in Microfabricated Electrohydrodynamic Systems," from Micro Electro Mechanical Systems conference in Travemunde Germany (1992).
- Hames et al., *Nuclear acid hybridization, a practical approach*, cover page and table of contents (1985).
- Hanahan et al., "Plasmid Screening at High Colony Density," *Meth. Enzymology*, 100:333-342 (1983).
- Hanahan et al., "Plasmid screening at high colony density," *Gene*, 10:63-67 (1980).
- Haridasan et al., "Peptide Synthesis using Photolytically Cleavable 2-Nitrobenzyloxycarbonyl Protecting Group," *Proc. Indian Natn. Sci. Acad.*, 53A(6):717-728 (1987).
- Harrison et al., "Capillary Electrophoresis and Sample Injection Systems Integrated on a Planar Glass Chip," *Anal. Chem.*, 64:1926-1932 (1992).
- Harrison et al., "Micromachining a Minaturized Capillary Electrophoresis-Based Chemical Analysis System on a Chip," *Science*, 261:895-897 (1993).
- Harrison et al., "Towards minaturized electrophoresis and chemical analysis systems on silicon: an alternative to chemical sensors*," *Sensors and Actuators*, B10:107-116 (1993).
- Harrison et al., "Rapid separation of fluorescein derivatives using a micromachined capillary electrophoresis system," *Analytica Chimica Acta*, 283:361-366 (1993).
- Hellberg et al., "Minimum analogue peptide sets (MAPS) for quantitative structure-activity relationships," *Int. J. Peptide Protein Res.*, 37:414-424 (1991).
- Hilser et al., "Protein and peptide mobility in capillary zone electrophoresis, A comparison of existing models and further analysis," *J. Chromatography*, 630:329-336 (1993).
- Ho et al., "Highly Stable Biosensor Using an Artificial Enzyme," *Anal.Chem.*, 59:536-537 (1987).
- Hochgeschwender et al., "Preferential expression of a defined T-cell receptor β -chain gene in hapten-specific cytotoxic T-cell clones," *Nature*, 322:376-378 (1986).
- Hodgson, J., "Assays A La Photolithography," *Biotech.*, 9:419 (1991).

US 6,440,667 B1

Page 7

- Hodgson et al., "Hybridization probe size control: optimized 'oligolabelling'," *Nuc. Acids Res.*, 15(15):6295 (1987).
- Hopman et al., "Bi-color detection of two target DNAs by non-radioactive in situ hybridization*," *Histochem.*, 85:1-4 (1986).
- Iwamura et al., "1-Pyrenylmethyl Esters, Photolabile Protecting Groups for Carboxylic Acids," *Tetrahedron Lett.*, 28(6):679-682 (1987).
- Iwamura et al., "1-(α -Diazobenzyl)pyrene: A Reagent for Photolabile and Fluorescent Protection of Carboxyl Groups of Amino Acids and Peptides," *Synlett*, p. 35-36 (1991).
- Jacobson et al., "Effects of Injection Schemes and Column Geometry on the Performance of Microchip Electrophoresis Devices," *Anal. Chem.*, 66:1107-1113 (1994).
- Jacobsen et al., "Open Channel Electrochromatography on a Microchip," *Anal. chem.*, 66:2369-2373 (1994).
- Jacobson et al., "Microchip Capillary Electrophoresis with an Integrated Postcolumn Reactor" *Anal. Chem.*, 66:3472-3476 (1994).
- Jacobson et al., "Precolumn Reactions with Electrophoretic Analysis Integrated on a Microchip," *Anal. Chem.*, 66:4127-4132 (1994).
- Jacobson et al., "Microfabricated chemical measurement systems," *Nature Medicine*, 1(10):1093-1096 (1995).
- Jacobsen et al., "Fused Quartz Substrates for Microchip Electrophoresis," *Anal. chem.*, 67:2059-2063 (1995).
- Jacobson et al., "High-Speed Separations on a Microchip," *Anal. Chem.*, 66:1114-1118 (1994).
- Jacobson et al., "Microchip electrophoresis with sample stacking," *Electrophoresis*, 16:481-486 (1995).
- Jayakumari, "Peptide synthesis in a triphasic medium catalysed by papain immobilized on a crosslinked polystyrene support," *Indian J. Chemistry*, 29B:514-517 (1990).
- Jovin et al., "Liminescence Digital Imaging Microscopy," *Ann. Rev. Biophys. Chem.*, 18:271-308 (1989).
- Kafatos et al., "Determination of nucleic acid sequence homologies and relative concentrations by a dot hybridization procedure," *Nuc. Acids Res.*, 7(6):1541-1553 (1979).
- Kaiser et al., "Peptide and Protein Synthesis by Segment Synthesis-Condensation," *Science*, 243:187-192 (1989).
- Kaplan et al., "Photolabile chelators for the rapid photorelease of divalent cations," *PNAS*, 85:6571-6575 (1988).
- Karube, "Micro-biosensors based on silicon fabrication technology," chapter 25 from *Biosensors: Fundamentals and Applications*, Turner et al., eds., Oxford Publ., 1987, pp. 471-480 (1987).
- Kates et al., "A Novel, Convenient, Three-dimensional Orthogonal Strategy for Solid-Phase Synthesis of Cyclic Peptides 1-3," *Tetrahed. Letters*, 34(10):1549-1552 (1993).
- Kerkof et al., "A Procedure for Making Simultaneous Determinations of the Relative Levels of Gene Transcripts in Tissues or Cells," *Anal. Biochem.*, 188:349-355 (1990).
- Khrapko et al., "An Oligonucleotide hybridization approach to DNA sequencing," *FEBS Lett.*, 256(1,2):118-122 (1989).
- Khrapko et al., "A method for DNA sequencing by hybridization with oligonucleotide matrix," *DNA Seq. Map.*, 1:375-388 (1991).
- Kidd et al., " α_1 -Antitrypsin deficiency detection by direct analysis of the mutation in the gene," *Nature*, 304:230-234 (1983).
- Kievits et al., "Rapid subchromosomal localization of cosmids by nonradioactive in situ hybridization," *Cytogenetics Cell Genetics*, 53(2-3):134-136 (1990).
- Kimura et al., "An Immobilized Enzyme Membrane Fabrication Method using an Ink Jet Nozzle," *Biosensors*, 4:41-52 (1988).
- Kimura et al., "An Integrated SOS/FET Multi-Biosensor," *Sensors & Actuators*, 9:373-387 (1986).
- Kitazawa et al., "In situ DNA-RNA hybridization using in vivo bromodeoxyuridine-labeled DNA probe," *Histochemistry*, 92:195-199 (1989).
- Kleinfeld et al., "Controlled Outgrowth of Dissociated Neurons on Patterned Substrates," *J. Neurosci.*, 8(11):4098-4120 (1988).
- Knight, P., "Materials and Methods/Microsequencers for Proteins and Oligosaccharides," *BioTech.*, 7:1075-76 (1989).
- Kohara et al., "The Physical Map of the Whole *E. coli* Chromosome: Application of a New Strategy for Rapid Analysis and Sorting of a Large Genomic Library," *Cell*, 50:495-508 (1987).
- Krile et al., "Multiplex holography with chirp-modulated binary phase-coded reference-beam masks," *Applied Opt.*, 18(1):52-56 (1979).
- Labat, I., "Subfragments as an informative characteristic of the DNA molecule—computer simulation," research report submitted to the University of Belgrade College of Natural Sciences and Mathematics, (1988).
- Lander et al., "Genomic Mapping by Fingerprinting Random Clones: A Mathematical Analysis," *Genomics*, 2:231-239 (1988).
- Lainer et al., "Human Lymphocyte Subpopulations Identified by Using Three-Color Immunofluorescence and Flow Cytometry Analysis: Correlation of Leu-2, Leu-3, Leu-7, Leu-8, and Leu-11 Clee Surface Antigen Expression," *Journal of Immunology*, 132(1):151-156 (1984).
- Lam et al., "A new type of synthetic peptide library for identifying ligand-binding activity," *Nature*, 354:82-84 (1991).
- Laskey et al., "Messenger RNA prevalence in sea urchin embryos measured with cloned cDNAs," *PNAS*, 77(9):5317-5321 (1980).
- Lee et al., "synthesis of a Polymer Surface Containing Covalently Attached Triethoxysilane Functionality: Adhesion to Glass," *Macromolecules*, 21:3353-3356 (1988).
- Lehrach et al., "Labelling oligonucleotides to high specific activity (I)," *Nuc. Acids Res.*, 17(12):4605-4610 (89).
- Lehrach et al., "Phage Vectors—EMBL Series," *Meth. Enzymology*, 153:103-115 (1987).
- Lehrach et al., "Hybridization Fingerprinting in Genome Mapping and Sequencing," *Genome Analysis vol. 1: Genetic and Physical Mapping*, Cold Spring Harbor Laboratory Press, pp. 39-81 (1990).
- Levy, M.F., "Preparing Additive Printed Circuits," *IBM Tech. Discl. Bull.*, 9(11):1473 (1967).
- Lewin, Benjamin, eds., *Genes*, third edition, John Wiley & Sons, cover page, preface and table of contents, (1987).
- Lichter et al., "High-Resolution Mapping of Human Chromosome 11 by in Situ hybridization with Cosmid Clones," *Science*, 247:64-69 (1990).
- Lichter et al., "Fluorescence in situ hybridization with Alu and L1 polymerase chain reaction probes for rapid characterization of human chromosomes in hybrid cell lines," *PNAS*, 87:6634-6638 (1990).
- Lichter et al., "Rapid detection of human chromosome 21 aberrations by in situ hybridization," *PNAS*, 85:9664-9668 (1988).

US 6,440,667 B1

Page 8

- Lichter et al., "Is non-isotopic in situ hybridization finally coming of age," *Nature*, 345:93-94 (1990).
- Lieberman et al., "A Light source Smaller Than the Optical Wavelength," *Science*, 247:59-61 (1990).
- Lipshutz et al., "Using Oligonucleotide Probe Arrays To Access Genetic Diversity," *BioTech.*, 19(3):442-7 (1995).
- Little, P., "Clone maps made simple," *Nature*, 346:611-612 (1990).
- Liu et al., "Sequential Injection Analysis in Capillary Format with an Electroosmotic Pump," *Talanta*, 41(11):1903-1910 (1994).
- Lockhart et al., "Expression monitoring by hybridization to high-density oligonucleotide arrays," *Nat. Biotech.*, 14:1675-1680 (1996).
- Logue et al., "General Approaches to Mask Design for Binary Optics," SPIE, 1052:19-24 (1989).
- Loken et al., "three-color Immunofluorescence Analysis of Leu Antigens on Human Peripheral Blood Using Two Lasers on a Fluorescence-Activated Cell Sorter," *Cytoetry*, 5:151-158 (1984).
- Love et al., "Screening of γ Library for Differentially Expressed Genes Using in Vitro Transcripts," *Anal. Biochem.*, 150:429-441 (1985).
- Lowe, C.R., "Biosensors," *Trends in Biotech.*, 2:59-65 (1984).
- Lowe, C.R., "An Introduction to the Concepts and Technology of Biosensors," *Biosensors*, 1:3-16 (1985).
- Lowe, C. R., *Biotechnology and Crop Improvement and Protection*, BCPC Publications, pp. 131-138 (1986).
- Lowe et al., "Solid-Phase Optoelectronic Biosensors," *Methods in Enzymology*, 137:338-347 (1988).
- Lowe, C.R., "Biosensors," *Phil. Tran. R. Soc. Lond.*, 324:487-496 (1989).
- Lu et al., "Differential screening of murine ascites cDNA libraries by means of in vitro transcripts of cell-cycle-phase-specific cDNA and digital image processing," *Gene*, 86:185-192 (1990).
- Luo, J. et al., "Improving the fidelity of *Thermus thermophilus* DNA ligase," *Nuc.Acids Res.*, 24(14):3071-3078 (1996).
- Lysov et al., "A new method for determining the DNA nucleotide sequence by hybridization with oligonucleotides," *Doklady Biochem.*, 303(1-6):436-438 (1989).
- Lysov et al., "DNA Sequencing by Oligonucleotide Hybridization," First International Conference on Electrophoresis, Supercomputing and the Human Genome, 4/10-13/90 p. 157.
- MacDonald et al., "A Rapid ELISA for Measuring Insulin in a Large Number of Research Samples," *Metabolism*, 38(5):450-452 (1989).
- Mairanovsky, V.G., "Electro-Deprotection-Electrochemical Removal of Protecting Groups*," *Agnew. Chem. Int. Ed. Engl.*, 15(5):281-292 (1976).
- Manz et al., "Miniaturized Total Chemical Analysis Systems: a Novel Concept for Chemical Sensing," *Sensors and Actuators*, B1:244-248 (1990).
- Manz et al., "Micromachining of monocrystalline silicon and glass for chemical analysis systems, A look into next century's technology or just a fashionable craze?," *Trends in Analytical Chem.*, 10(5):144-149 (1991).
- Manz et al., "Planar chips technology for minaturization and integration of separation techniques into monitoring systems, Capillary electrophoresis on a chip," *J. Chromatography*, 593:253-258 (1992).
- Manz et al., "Planar Chips Technology for Miniaturization of Separation Systems: A Developing Perspective in Chemical Monitoring," chapter 1, 1-64 (1993).
- Manz et al., "Electroosmotic pumping and electrophoretic separations for miniaturized chemical analysis systems," *J. Micromech. Microeng.*, 4:257-265 (1994).
- Masiakowski et al., "Cloning of cDNA sequences of hormone-regulated genes from the MCF-7 human breast cancer cell line," *Nuc. Acids Res.*, 10(24):7895-7903 (1982).
- Matsumoto et al., "Preliminary Investigation of Micropumping Based on Electrical Control of Interfacial Tension," *IEEE*, pp. 105-110 (1990).
- Matsuzawa et al., "Containment and growth of neuroblastoma cells on chemically patterned substrates," *J. Neurosci. Meth.*, 50:253-260 (1993).
- Matthes et al., "Simultaneous rapid chemical synthesis of over one hundred oligonucleotides on a microscale," *EMBO J.*, 3(4):801-805 (1984).
- McCray et al., "Properties and Uses of Photoreactive Caged Compounds," *Ann. Rev. Biophys. Biophys. Chem.*, 18:239-270 (1989).
- McGall et al., "The Efficiency of Light-Directed Synthesis of DNA Arrays on Glass Substrates," *J. American Chem. Soc.*, 119(22):5081-5090 (1997).
- McGillis, VLSI Technology, Sze, eds., Chapter 7, "Lithography," pp. 267-301 (1983).
- McMurray, J.S., "Solid Phase Synthesis of a Cyclic Peptide Using Fmoc Chemistry," *Tetrahedron Letters*, 32(52):7679-7682 (1991).
- Meinkoth et al., "Review: Hybridization of Nucleic Acids Immobilized on solid Supports," *Analytical Biochem.*, 138:267-284 (1984).
- Melcher et al., "Traveling-Wave Bulk Electroconvection Induced across a Temperature Gradient," *Physics of Fluids*, 10(6):1178-1185 (1967).
- Merrifield, R.B., "Solid Phase peptide Synthesis. I. The Synthesis of a Tetrapeptide," *J.Am.Chem.Soc.*, 85:2149-2154 (1963).
- Michiels et al., "Molecular approaches to genome analysis: a strategy for the construction of ordered overlapping clone libraries," *CABIOS*, 3(3):203-10 (1987).
- Mirzabekov, A.D., "DNA sequencing by hybridization—a megasequencing method and a diagnostic tool?," *TIBTECH*, 12:27-32 (1994).
- Miyada et al., "Oligonucleotide Hybridization Techniques," *Meth. Enzymology*, 154:94-107 (1987).
- Monaco et al., "Human Genome Linking with Cosmids and Yeast Artificial Chromosomes", abstract from CSHS, p. 50, (1989).
- Morita et al., "Direct pattern fabrication on silicone resin by vapor phase electron beam polymerization," *J.Vac.Sci.Technol.*, B1(4):1171-1173 (1983).
- Morrison et al., "Solution-Phase Detection of Polynucleotides Using Interacting Fluorescent Labels and Competitive Hybridization," *Anal. Biochem.*, 183:231-244 (1989).
- Munegumi et al., "thermal Synthesis of Polypeptides from N-Boc-Amino Acid (Aspartic Acid, β -Aminoglutaric Acid) Anhydrides," *Chem. Letters*, pp. 1643-1646 (1988).
- Mutter et al., "Impact of Conformation on the Synthetic Strategies for Peptide Sequences," pp. 217-228 from *Chemistry of Peptides and Proteins*, vol. 1, Proceedings of the Third USSR-FRG Symp., in USSR (1982).

US 6,440,667 B1

Page 9

- Nakamori et al., "A Simple and Useful Method for Simultaneous Screening of Elevated Levels of Expression of a Variety of Oncogenes in Malignant Cells," *Jpn. J. Cancer Res.*, 79:1311-1317 (1988).
- Nederlof et al., "Multiple Fluorescence In Situ Hybridization," *Cytometry*, 11:126-131 (1990).
- Nederlof et al., "Three-Color Fluorescence In Situ Hybridization for the Simultaneous Detection of Multiple Nucleic Acid Sequences," *Cytometry*, 10:20-27 (1989).
- Nizetic et al., "An improved bacterial colony lysis procedure enables direct DNA hybridisation using short (10, 11 bases) oligonucleotides to cosmids," *Nuc. Acids Res.*, 19(1):182 (1990).
- Nizetic et al., "Construction, arraying, and high-density screening of large insert libraries of human chromosomes X and 21: their potential use as reference libraries," *PNAS*, 88:3233-3237 (1991).
- Nyborg, W., "Acoustic Streaming," chapter 11 pp. 265-329 from *Physical Acoustics, Principles and Methods*, Mason, eds., vol. II, part B, Academic Press, New York and London (1965).
- Oevirk et al., "High Performance Liquid Chromatography Partially Integrated onto a Silicon Chip," *Analyt. Meth. Instrumentation*, 2(2):74-82 (1995).
- Ohtsuka et al., "Studies on transfer ribonucleic acids and related compounds. IX Ribonucleic oligonucleotide synthesis using a photosensitive 0-nitrobenzyl protection at the 2'-hydroxyl group," *Nuc.Acids.Res.*, 1(10):1351-1357 (1974).
- Olefirowicz et al., "Capillary Electrophoresis for Sampling Single Nerve Cells," *Chimia*, 45(4):106-108 (1991).
- Olson et al., "Random-clone strategy for genomic restriction mapping in yeast," *PNAS*, 83:7826-7830 (1986).
- Patchornik et al., "Photosensitive Protecting Groups," *J.Am.Chem.Soc.*, 92(21):6333-6335 (1970).
- Patent Abstracts of Japan from EPO, Abst. 13:557, JP 1-233 447 (1989).
- Pease et al., "Light-generated oligonucleotide arrays for rapid DNA sequence analysis," *PNAS*, 91:5022-26 (1994).
- Pevzner, P.A., "DNA Physical Mapping and Alternating Eulerian Cycles in Colored Grapes," *Algorithmica*, 13(1-2):77-105 (1995).
- Pevzner et al., "Multiple Filtration and Approximate Pattern Matching," *Algorithmica*, 13(1-2):135-154 (1995).
- Pevzner et al., "Generalized Sequence Alignment and Duality," *Adv. Applied Math.*, 14:139-171 (1993).
- Pevzner, P.A., "1-Tuple DNA Sequencing: Computer Analysis," *J. Biomol. Struct. Dynam.*, 7(1):63-69 (1989).
- Pfahler et al., "Liquid Transport in Micron and Submicron Channels," *Sensors and Actuators*, A21-A23:431-4 (90).
- Pfeifer et al., "Genomic Sequencing and Methylation Analysis by Ligation Mediated PCR," *Science*, 246:810-813 (1989).
- Pidgeon et al., "Immobilized Artificial Membrane Chromatography: Supports Composed of Membrane Lipids," *Anal. Biochem.*, 176:36-47 (89).
- Pillai, V.N., "Photoremovable Protecting Groups in Organic Synthesis," *Synthesis*, pp. 1-26 (1980).
- Pillai et al., "3-Nitro-4-Aminomethylbenzoyl derivative von Polyethylenglykolen: Eine neue Klasse von Photosensitiven loslichen Polymeren Trägern zur Synthese von C-terminalen Peptidamiden," *Tetrah. Ltr.*, #36 p. 3409-3412 (1979).
- Pillai et al., "Synthetic Hydrophilic Polymers, Biomedical and Chemical Applications," *Naturwissenschaften*, 68:558-566 (1981).
- Pirrung et al., "Proofing of Photolithographic DNA Synthesis with 3'.5'-Dimethoxybenzoinyloxycarbonyl-Protected Deoxynucleoside Phosphoramidites," *J. Org. Chem.*, 63(2):241-246 (1998).
- Pirrung et al., "Comparison of Methods for Photochemical Phosphoramidite-Based DNA Synthesis," *J. Org. Chem.*, 60:6270-6276 (1995).
- Ploax et al., "Cyclization of peptides on a solid support," *Int. J. Peptide Protein Research*, 29:162-169 (1987).
- Polsky-Cynkin et al., "Use of DNA Immobilized on Plastic and Agarose Supports to Detect DNA by Sandwich Hybridization," *Clin. Chem.*, 31(9):1428-1443 (1985).
- Poustka et al., "Molecular Approaches to Mammalian Genetics," Cold Spring Harbor Symposia on Quantitative Biology, 51:131-139 (1986).
- Purushothaman et al., "Synthesis of 4,5-diarylimidazole-2-thiones and their photoconversion to bis(4,5-diarylimidazole-2-yl) sulphides," *Ind. J. Chem.*, 29B:18-21 (1990).
- Quesada et al., "High-Sensitivity DNA Detection with a Laser-Excited Confocal Fluorescence Gel Scanner," *Biotechniques*, 10:616 (1991).
- Reichmanis et al., *J. Polymer Sci. Polymer Chem. Edition*, 23:1-8 (1985).
- Renz et al., "A colorimetric method for DNA hybridization," *Nuc. Acids Res.*, 12(8):3435-3445 (1984).
- Richter et al., "An Electrohydrodynamic Micropump," *IEEE*, pp. 99-104 (1990).
- Richter et al., "Electrohydrodynamic Pumping and Flow Measurement," *IEEE*, pp. 271-276 (1991).
- Richter et al., "A Micromachined electrohydrodynamic (EHD) pump," *Sensors and Actuators*, A29:159-168 (91).
- Robertson et al., "A General and Efficient Route for Chemical Aminoacylation of Transfer RNAs," *J. Am. Chem. Soc.*, 113:2722-2729 (1991).
- Rodda et al., "The Antibody Response to Myoglobin-I. Systematic Synthesis of Myoglobin Peptides Reveals Location and Substructure of Species-Dependent Continuous Antigenic Determinants," *Mol. Immunol.*, 23(6):603-610 (1986).
- Rodgers, R.P., "Data Processing of Immunoassay Results," *Manual of Clin. Lab. Immunol.*, 3rd ed., ch. 15, pp. 82-87 (1986).
- Rose, D.J., "Free-solution reactor for post-column fluorescence detection in capillary zone electrophoresis," *J. Chromatography*, 540:343-353 (1991).
- Rovero et al., "Synthesis of Cyclic Peptides on solid Support," *Tetrahed. Letters*, 32(23):2639-2642 (1991).
- Sambrook, *Molecular Cloning—A Laboratory Manual*, publ. in 1989 (no included).
- Saiki et al., "Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes," *PNAS*, 86:6230-6234 (1989).
- Saiki et al., "Analysis of enzymatically amplified β -globin and HLA-DQ α DNA with Allele-specific oligonucleotide probes," *Nature*, 324:163-166 (1986).
- Schafer et al., "DNA fingerprinting using non-radioactive oligonucleotide probes specific for simple repeats," *Nuc. Acids Res.*, 16(19):9344 (1988).

US 6,440,667 B1

Page 10

- Scharf et al., "HLA class II allelic variation and susceptibility to pemphigus vulgaris," *PNAS*, 85(10):3504-3508 (1988).
- Schena et al., "Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes," *PNAS*, 93:10614-10619 (1996).
- Schuup et al., "Mechanistic Studies of the Photorearrangement of o-Nitrobenzyl Esters," *J. Photochem.*, 36:85-97 (1987).
- Seed, B., "Diazotizable arylamine cellulose papers for the coupling and hybridization of nucleic acids," *Nuc. Acids Res.*, 10(5):1799-1810 (1982).
- Seiler et al., "Planar Glass Chips for Capillary Electrophoresis: Repetitive Sample Injection, Quantitation, and Separation Efficiency," *Anal. Chem.*, 65:1481-1488 (1993).
- Seller et al., "Electroosmotic Pumping and Valveless Control of Fluid Flow within a Manifold of Capillaries on a Glass Chip," *Anal. Chem.*, 66:3485-3491 (1994).
- Semmelhack et al., "Selective Removal of Protecting Groups Using Controlled Potential Electrolysis," *J. Am. Chem. Society*, 94(14):5139-5140 (1972).
- Sheldon et al., "Matrix DNA Hybridization," *Clinical Chemistry*, 39(4):718-719 (1993).
- Shin et al., "Dehydrooligonopeptides. XI. Facile Synthesis of Various Kinds of Dehydrodi- and tripeptides, and Dehydroenkephalins Containing Tyr Residue by Using N-Carboxydehydrotyrosine Anhydride," *Bull. Chem. Soc. Jpn.*, 62:1127-1135 (1989).
- Sim et al., "Use of a cDNA Library for Studies on Evolution and Developmental Expression of the Chorion Multigene Families," *Cell*, 18:1303-1316 (1979).
- Smith et al., "A Novel Method for Delineating Antigenic Determinants: Peptide Synthesis and Radioimmunoassay Using the Same Solid Support," *Immunochemistry*, 14:565-568 (1977).
- Sofia, M.J., "Carbohydrate-based combinatorial libraries," *Molecular Diversity*, 3:75-94 (1998).
- Southern et al., "Report on the Sequencing by Hybridization Workshop," *Genomics*, 13:1378-1383 (1992).
- Southern et al., "Oligonucleotide hybridisations on glass supports: a novel linker for oligonucleotide synthesis and hybridization properties of oligonucleotides synthesized in situ," *Nuc. Acids Res.*, 20(7):1679-1684 (1992).
- Southern et al., "Analyzing and Comparing Nucleic Acid Sequences by Hybridization to Arrays of Oligonucleotides: Evaluation Using Experimental Models," *Genomics*, 13:1008-10017 (1992).
- Southern, E.M., "Detection of Specific Sequences Among DNA Fragments Separated by Gel Electrophoresis," *J. Mol. Biol.*, 98:503-517 (1975).
- Stemme et al., "A valveless diffuser/nozzle-based fluid pump," *Sensors and Actuators*, A39:159-167 (1993).
- Stryer, L., "DNA Probes and Genes Can be Synthesized by Automated Solid-Phase Methods," from *Biochemistry*, Third Edition, published by W.H. Freeman & Co., (1988).
- Stuber et al., "Synthesis and photolytic cleavage of bovine insulin B22-30 on a nitrobenzoylglycyl-poly (ethylene glycol) support," *Int. J. Peptide Protein Res.*, 22(3):277-283 (1984).
- Sundberg et al., "Spatially-Addressable Immobilization of Macromolecules on Solid Supports," *J. Am. Chem. Soc.*, 117(49):12050-12057 (1995).
- Swedberg, S.A., "Use of non-ionic and zwitterionic surfactants to enhance selectivity in high-performance capillary electrophoresis, An apparent micellar electrokinetic capillary chromatography mechanism," *J. Chromatography*, 503:449-452 (1990).
- Thomas, P.S., "Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose," *PNAS*, 77(9):5201-5205 (1980).
- Titus et al., "Texas Red, a Hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow microfluorometric and fluorescence microscopic studies," *J. Immunol. Meth.*, 50:193-204 (1982).
- Tkachuk et al., "Detection of bcr-abl Fusion in chronic Myelogenous Leukemia by in situ Hybridization," *Science*, 250:559-562 (90).
- Trzeciak et al., "Synthesis of 'Head-to-Tail' Cyclized Peptides on Solid Support by Fmoc Chemistry," *Tetrahed. Letters*, 33(32):4557-4560 (1992).
- Tsien et al., "Control of Cytoplasmic Calcium with Photolabile Tetracarboxylate 2-Nitrobenzhydryl Chelators," *Biophys. J.*, 50:843-853 (1986).
- Tsutsumi et al., "Expression of L- and M-Type Pyruvate Kinase in Human Tissues," *Genomics*, 2:86-89 (1988).
- Turchinskii et al., "Multiple Hybridization in Genome Analysis, Reaction of Diamines and Bisulfate with Cytosine for Introduction of Nonradioactive labels Into DNA," *Molecular Biology*, 22:1229-1235 (1988).
- Turner et al., "Photochemical Activation of Acylated α -Thrombin," *J. Am. Chem. Soc.*, 109:1274-1275 (1987).
- Urdea et al., "A novel method for the rapid detection of specific nucleotide sequences in crude biological samples without blotting or radioactivity; application to the analysis of hepatitis B virus in human serum," *Gene*, 61:253-264 (1987).
- Urdea et al., "A comparison of non-radioisotopic hybridization assay methods using fluorescent, chemiluminescent and enzyme labeled synthetic oligodeoxyribonucleotide probes," *Nuc. Acids Res.*, 16(11):4937-4956 (1988).
- Van der Voort et al., "Design and Use of a Computer Controlled Confocal Microscope for Biological Applications," *Scanning*, 7(2):66-78 (1985).
- Van Hijfte et al., "Intramolecular 1,3-Diyl Trapping Reactions. A Formal Total Synthesis of -Coriolin," *J. Organic Chemistry*, 50:3942-3944 (1985).
- Veldkamp, W.B., "Binary optics: the optics technology of the 1990s," *CLEO 90*, vol. 7, paper # CMG6 (1990).
- Verlaan-de Vries et al., "A dot-blot screening procedure for mutated ras oncogenes using synthetic oligodeoxynucleotides," *Gene*, 50:313-320 (1986).
- Verpoorte et al., "Three-dimensional micro flow manifolds for miniaturized chemical analysis systems," *J. Micromech. Microeng.*, 4:246-256 (1994).
- Volkmut et al., "DNA electrophoresis in microlithographic arrays," *Nature*, 358:600-602 (1992).
- Voss et al., "The immobilization of oligonucleotides and their hybridization properties," *Biochem. Soc. Transact.*, 16:216-217 (1988).
- Wada, A., *International Workshop on Automatic and High Speed DNA Base Sequencing*, Hayashibara Forum 1987 at Hayashibara Biochemical Laboratories, Okayama, Japan, Jul. 7-9, 1987.

US 6,440,667 B1

Page 11

- Walker et al., "Photolabile Protecting Groups for an Acetylcholine Receptor Ligand. Synthesis and Photochemistry of a New Class of o-Nitrobenzyl Derivatives and their Effects on Receptor Function," *Biochemistry*, 25:1799-1805 (1986).
- Wallace et al., "The use of synthetic oligonucleotides as hybridization probes. II. Hybridization of oligonucleotides of mixed sequence to rabbit β -globin DNA," *Nuc. Acids Res.*, 9(4):879 (1981).
- Wallace et al., "Hybridization of synthetic oligodeoxyribonucleotides to $\Phi\chi$ 174 DNA: the effect of single base pair mismatch," *Nuc. Acids Res.*, 11(6):3543-3557 (1979).
- Washizu et al., "Handling Biological Cells Using a Fluid Integrated Circuit," *IEEE Transactions Industry Applications*, 26(2):352-358 (1990).
- Wiedmann, M. et al., "Ligase Chain Reaction (LCR)—Overview and Applications," *PCR Meth. Appl.*, 3(4):S51-S64 (1994).
- Werner et al., "Size-Dependent Separation of Proteins Denatured in SDS by Capillary Electrophoresis Using a Replaceable Sieving Matrix," *Anal. Biochem.*, 212:253-258 (1993).
- White et al., "An Evaluation of Confocal Versus Conventional Imaging of Biological Structures by Fluorescence Light Microscopy," *J. Cell Biol.*, 105(1):41-48 (1987).
- Widacki et al., "Biochemical Differences in Qa-2 Antigens Expressed by Qa-2+,6+ and Qa-2a+,6- Strains. Evidence for Differential Expression of the Q7 and Q9 Genes," *Mol. Immunology*, 27(6):559-570 (1990).
- Wilcox et al., "Synthesis of Photolabile 'Precursors' of Amino Acid Neurotransmitters," *J. Org. Chem.*, 55:1585-1589 (1990).
- Wilding et al., "PCR in a Silicon Microstructure," *Clin. Chem.*, 40(9):1815-1818 (1994).
- Wilding et al., "Manipulation and Flow of Biological Fluids in Straight Channels Micromachined in Silicon," *Clin. Chem.*, 40(1):43-47 (1994).
- Wittman-Liebold, eds., *Methods in Protein Sequence Analysis*, from Proceedings of 7th Int'l Conf., Berlin, Germany, Jul. 3-8, 1988, table of contents, pp. xi-xx* (1989).
- Wood et al., "Base composition-independent hybridization in tetramethylammonium chloride: A method for oligonucleotide screening of highly complex gene libraries," *PNAS*, 82:1585-1588 (1985).
- Woolley et al., "Ultra-high-speed DNA fragment separations using microfabricated capillary array electrophoresis chips," *PNAS*, 91:11348-11352 (1994).
- Wu et al., "Synthesis and Properties of Adenosine-5'-triphospho- γ -5-(5-sulfonic acid)naphthyl Ethylamide: A Fluorescent Nucleotide Substrate for DNA-Dependent RNA Polymerase from *Escherichia coli*," *Arch. Biochem. Biophys.*, 246(2):564-571 (1986).
- Wu et al., "Laboratory Methods, Direct Analysis of Single Nucleotide Variation in Human DNA and RNA Using In Situ Dot Hybridization," *DNA*, 8(2):135-142 (1989).
- Yamamoto et al., "Features and applications of the laser scanning microscope," *J. Mod. Optics*, 37(11):1691-1701 (1990).
- Yarbrough et al., "Synthesis and Properties of Fluorescent Nucleotide Substrates for DNA-dependent RNA Polymerases," *J. Biol. Chem.*, 254(23):12069-12073 (1979).
- Yosomiya et al., "Performance, Glass fiber Having Isocyanate Group on the Surface. Preparation and Reaction with Amino Acid," *Polymer Bulletin*, 12:41-48 (1984).
- Young, W.S., "Simultaneous Use of Digoxigenin- and Radiolabeled Oligodeoxyribonucleotide Probes for Hybridization Histochemistry," *Neuropeptides*, 13:271-275 (1989).
- Yue et al., "Miniature Field-Flow Fractionation System for Analysis of Blood Cells," *Clin. Chem.*, 40(9):1810-1814 (1994).
- Zehavi et al., "Light-Sensitive Glycosides. I. 6-Nitroveratryl β -D-Glucopyranoside and 2-Nitrobenzyl β -D-Glucopyranoside," *J. Org. Chem.*, 37(14):2281-2285 (1972).
- Zengerle et al., "Transient measurements on miniaturized diaphragm pumps in microfluid systems," *Sensors and Actuators*, A46-47:557-561 (1995).
- Zischler et al., "Non-radioactive oligonucleotide fingerprinting in the gel," *Nuc. Acids Res.*, 17(11):4411 (1989).
- Zischler et al., "Digoxigenated oligonucleotide probes specific for simple repeats in DAN fingerprinting and hybridization in situ," *Hum. Genet.*, 82:227-233 (1989).

* cited by examiner




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 2G




FIG. 2H




FIG. 2I




FIG. 2K




FIG. 2L




FIG. 2M




FIG. 2N

US 6,440,667 B1

1

ANALYSIS OF TARGET MOLECULES USING AN ENCODING SYSTEM

CROSS REFERENCE TO RELATED APPLICATIONS

This is a divisional of Appln. Ser. No. 09/056,927, filed Apr. 8, 1998 now U.S. Pat. No. 6,197,506, which is a continuation of Appln. Ser. No. 08/670,118, filed Jun. 25, 1996, now U.S. Pat. No. 5,800,992; which is a divisional of Appln. Ser. No. 08/168,904, filed Dec. 15, 1993, now abandoned; which is a continuation of Appln. Ser. No. 07/624,114 filed Dec. 6, 1990, now abandoned; which is a continuation in-part of commonly assigned Appln. Ser. No. 07/492,462, filed Mar. 7, 1990, now U.S. Pat. No. 5,143,854; and Appln. Ser. No. 07/362,901, filed Jun. 7, 1989, now abandoned which are hereby incorporated by reference.

Additional commonly assigned Appln. Ser. Nos. 07/624,120 and 07/626,730, both of which were filed on Dec. 6, 1990; Appln. Ser. No. 07/435,316, filed Nov. 13, 1989, now abandoned; and U.S. Pat. No. 5,252,743 are also hereby incorporated herein by reference.

BACKGROUND OF THE INVENTION

The present invention relates to the sequencing, fingerprinting, and mapping of polymers, particularly biological polymers. The inventions may be applied, for example, in the sequencing, fingerprinting, or mapping of nucleic acids, polypeptides, oligosaccharides, and synthetic polymers.

The relationship between structure and function of macromolecules is of fundamental importance in the understanding of biological systems. These relationships are important to understanding, for example, the functions of enzymes, structural proteins, and signalling proteins, ways in which cells communicate with each other, as well as mechanisms of cellular control and metabolic feedback.

Genetic information is critical in continuation of life processes. Life is substantially informationally based and its genetic content controls the growth and reproduction of the organism and its complements. Polypeptides, which are critical features of all living systems, are encoded by the genetic material of the cell. In particular, the properties of enzymes, functional proteins, and structural proteins are determined by the sequence of amino acids which make them up. As structure and function are integrally related, many biological functions may be explained by elucidating the underlying structural features which provide those functions. For this reason, it has become very important to determine the genetic sequences of nucleotides which encode the enzymes, structural proteins, and other effectors of biological functions. In addition to segments of nucleotides which encode polypeptides, there are many nucleotide sequences which are involved in control and regulation of gene expression.

The human genome project is directed toward determining the complete sequence of the genome of the human organism. Although such a sequence would not correspond to the sequence of any specific individual, it would provide significant information as to the general organization and specific sequences contained within segments from particular individuals. It would also provide mapping information which is very useful for further detailed studies. However, the need for highly rapid, accurate, and inexpensive sequencing technology is nowhere more apparent than in a demanding sequencing project such as this. To complete the sequencing of a human genome would require the determination of approximately 3×10^9 , or 3 billion base pairs.

2

The procedures typically used today for sequencing include the Sanger dideoxy method, see, e.g., Sanger et al. (1977) *Proc. Natl. Acad. Sci. USA*, 74:5463-5467, or the Maxam and Gilbert method, see, e.g., Maxam et al., (1980) *Methods in Enzymology*, 65:499-559. The Sanger method utilizes enzymatic elongation procedures with chain terminating nucleotides. The Maxam and Gilbert method uses chemical reactions exhibiting specificity of reaction to generate nucleotide specific cleavages. Both methods require a practitioner to perform a large number of complex manual manipulations. These manipulations usually require isolating homogeneous DNA fragments, elaborate and tedious preparing of samples, preparing a separating gel, applying samples to the gel, electrophoresing the samples into this gel, working up the finished gel, and analyzing the results of the procedure.

Thus, a less expensive, highly reliable, and labor efficient means for sequencing biological macromolecules is needed. A substantial reduction in cost and increase in speed of nucleotide sequencing would be very much welcomed. In particular, an automated system would improve the reproducibility and accuracy of procedures. The present invention satisfies these and other needs.

SUMMARY OF THE INVENTION

The present invention provides improved methods useful for de novo sequencing of an unknown polymer sequence, for verification of known sequences, for fingerprinting polymers, and for mapping homologous segments within a sequence. By reducing the number of manual manipulations required and automating most of the steps, the speed, accuracy, and reliability of these procedures are greatly enhanced.

The production of a substrate having a matrix of positionally defined regions with attached reagents exhibiting known recognition specificity can be used for the sequence analysis of a polymer. Although most directly applicable to sequencing, the present invention is also applicable to fingerprinting, mapping, and general screening of specific interactions. The VLSIPS™ Technology (Very Large Scale Immobilized Polymer Synthesis) substrates will be applied to evaluating other polymers, e.g., carbohydrates, polypeptides, hydrocarbon synthetic polymers, and the like. For these non-polynucleotides, the sequence specific reagents will usually be antibodies specific for a particular subunit sequence.

According to one aspect of the masking technique, the invention provides an ordered method for forming a plurality of polymer sequences by sequential addition of reagents comprising the step of serially protecting and deprotecting portions of the plurality of polymer sequences for addition of other portions of the polymer sequences using a binary synthesis strategy.

The present invention also provides a means to automate sequencing manipulations. The automation of the substrate production method and of the scan and analysis steps minimizes the need for human intervention. This simplifies the tasks and promotes reproducibility.

The present invention provides a composition comprising a plurality of positionally distinguishable sequence specific reagents attached to a solid substrate, which reagents are capable of specifically binding to a predetermined subunit sequence of a preselected multi-subunit length having at least three subunits, said reagents representing substantially all possible sequences of said preselected length. In some embodiments, the subunit sequence is a polynucleotide or a

US 6,440,667 B1

3

polypeptide, in others the preselected multi-subunit length is five subunits and the subunit sequence is a polynucleotide sequence. In other embodiments, the specific reagent is an oligonucleotide of at least about five nucleotides. Alternatively, the specific reagent is a monoclonal antibody. Usually the specific reagents are all attached to a single solid substrate, and the reagents comprise about 3000 different sequences. In other embodiments, the reagents represents at least about 25% of the possible subsequences of said preselected length. Usually, the reagents are localized in regions of the substrate having a density of at least 25 regions per square centimeter, and often the substrate has a surface area of less than about 4 square centimeters.

The present invention also provides methods for analyzing a sequence of a polynucleotide or a polypeptide, said method comprising the step of:

- a) exposing said polynucleotide or polypeptide to a composition as described.

It also provides useful methods for identifying or comparing a target sequence with a reference, said method comprising the step of:

- a) exposing said target sequence to a composition as described;
- b) determining the pattern of positions of the reagents which specifically interact with the target sequence; and
- c) comparing the pattern with the pattern exhibited by the reference when exposed to the composition.

The present invention also provides methods for sequencing a segment of a polynucleotide comprising the steps of:

- a) combining:
 - i) a substrate comprising a plurality of chemically synthesized and positionally distinguishable oligonucleotides capable of recognizing defined oligonucleotide sequences; and
 - ii) a target polynucleotide; thereby forming high fidelity matched duplex structures of complementary subsequences of known sequence; and
- b) determining which of said reagents have specifically interacted with subsequences in said target polynucleotide.

In one embodiment, the segment is substantially the entire length of said polynucleotide.

The invention also provides methods for sequencing a polymer, said method comprising the steps of:

- a) preparing a plurality of reagents which each specifically bind to a subsequence of preselected length;
- b) positionally attaching each of said reagents to one or more solid phase substrates, thereby producing substrates of positionally definable sequence specific probes;
- c) combining said substrates with a target polymer whose sequence is to be determined; and
- d) determining which of said reagents have specifically interacted with subsequences in said target polymer.

In one embodiment, the substrates are beads. Preferably, the plurality of reagents comprise substantially all possible subsequences of said preselected length found in said target. In another embodiment, the solid phase substrate is a single substrate having attached thereto reagents recognizing substantially all possible subsequences of preselected length found in said target.

In another embodiment, the method further comprises the step of analyzing a plurality of said recognized subsequences to assemble a sequence of said target polymer. In a

4

bead embodiment, at least some of the plurality of substrates have one subsequence specific reagent attached thereto, and the substrates are coded to indicate the sequence specificity of said reagent.

The present invention also embraces a method of using a fluorescent nucleotide to detect interactions with oligonucleotide probes of known sequence, said method comprising:

- a) attaching said nucleotide to a target unknown polynucleotide sequence, and
- b) exposing said target polynucleotide sequence to a collection of positionally defined oligonucleotide probes of known sequences to determine the sequences of said probes which interact with said target.

In a further refinement, an additional step is included of:

- a) collating said known sequences to determine the overlaps of said known sequences to determine the sequence of said target sequence.

A method of mapping a plurality of sequences relative to one another is also provided, the method comprising:

- a) preparing a substrate having a plurality of positionally attached sequence specific probes;
- b) exposing each of said sequences to said substrate, thereby determining the patterns of interaction between said sequence specific probes and said sequences; and
- c) determining the relative locations of said sequence specific probe interactions on said sequences to determine the overlaps and order of said sequences.

In one refinement, the sequence specific probes are oligonucleotides, applicable to where the target sequences are nucleic acid sequences.

In the nucleic acid sequencing application, the steps of the sequencing process comprise:

- a) producing a matrix substrate having known positionally defined regions of known sequence specific oligonucleotide probes;
- b) hybridizing a target polynucleotide to the positions on the matrix so that each of the positions which contain oligonucleotide probes complementary to a sequence on the target hybridize to the target molecule;
- c) detecting which positions have bound the target, thereby determining sequences which are found on the target; and
- d) analyzing the known sequences contained in the target to determine sequence overlaps and assembling the sequence of the target therefrom.

The enablement of the sequencing process by hybridization is based in large part upon the ability to synthesize a large number (e.g., to virtually saturate) of the possible overlapping sequence segments and distinguishing those probes which hybridize with fidelity from those which have mismatched bases, and to analyze a highly complex pattern of hybridization results to determine the overlap regions.

The detecting of the positions which bind the target sequence would typically be through a fluorescent label on the target. Although a fluorescent label is probably most convenient, other sorts of labels, e.g., radioactive, enzyme linked, optically detectable, or spectroscopic labels may be used. Because the oligonucleotide probes are positionally defined, the location of the hybridized duplex will directly translate to the sequences which hybridize. Thus, analysis of the positions provides a collection of subsequences found within the target sequence. These subsequences are matched with respect to their overlaps so as to assemble an intact target sequence.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates a flow chart for sequence, fingerprint, or mapping analysis.

US 6,440,667 B1

5

FIGS. 2A–2M illustrate the proper function of a VLSIPS™ Technology nucleotide synthesis.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Overall Description

- A. general
- B. VLSIPS substrates
- C. binary masking
- D. applications
- E. detection methods and apparatus
- F. data analysis

II. Theoretical Analysis

- A. simple n-mer structure; theory
- B. complications
- C. non-polynucleotide embodiments

III. Polynucleotide Sequencing

- A. preparation of substrate matrix
- B. labeling target polynucleotide
- C. hybridization conditions
- D. detection; VLSIPS scanning
- E. analysis
- F. substrate reuse
- G. non-polynucleotide aspects

IV. Fingerprinting

- A. general
- B. preparation of substrate matrix
- C. labeling target nucleotides
- D. hybridization conditions
- E. detection; VLSIPS scanning
- F. analysis
- G. substrate reuse
- H. non-polynucleotide aspects

V. Mapping

- A. general
- B. preparation of substrate matrix
- C. labeling
- D. hybridization/specific interaction
- E. detection
- F. analysis
- G. substrate reuse
- H. non-polynucleotide aspects

VI. Additional Screening

- A. specific interactions
- B. sequence comparisons
- C. categorizations
- D. statistical correlations

VII. Formation of Substrate

- A. instrumentation
- B. binary masking
- C. synthetic methods
- D. surface immobilization

VIII. Hybridization/Specific Interaction

- A. general
- B. important parameters

IX. Detection Methods

- A. labeling techniques
- B. scanning system

6

X. Data Analysis

- A. general
- B. hardware
- C. software

XI. Substrate Reuse

- A. removal of label
- B. storage and preservation
- C. processes to avoid degradation of oligomers

XII. Integrated Sequencing Strategy

- A. initial mapping strategy
- B. selection of smaller clones
- C. actual sequencing procedures

XIII. Commercial Applications

- A. sequencing
- B. fingerprinting
- C. mapping

I. OVERALL DESCRIPTION

A. General

The present invention relies in part on the ability to synthesize or attach specific recognition reagents at known locations on a substrate, typically a single substrate. In particular, the present invention provides the ability to prepare a substrate having a very high density matrix pattern of positionally defined specific recognition reagents. The reagents are capable of interacting with their specific targets while attached to the substrate, e.g., solid phase interactions, and by appropriate labeling of these targets, the sites of the interactions between the target and the specific reagents may be derived. Because the reagents are positionally defined, the sites of the interactions will define the specificity of each interaction. As a result, a map of the patterns of interactions with specific reagents on the substrate is convertible into information on the specific interactions taking place, e.g., the recognized features. Where the specific reagents recognize a large number of possible features, this system allows the determination of the combination of specific interactions which exist on the target molecule. Where the number of features is sufficiently large, the identical same combination, or pattern, of features is sufficiently unlikely that a particular target molecule may often be uniquely defined by its features. In the extreme, the features may actually be the subunit sequence of the target molecule, and a given target sequence may be uniquely defined by its combination of features.

In particular, the methodology is applicable to sequencing polynucleotides. The specific sequence recognition reagents will typically be oligonucleotide probes which hybridize with specificity to subsequences found on the target sequence. A sufficiently large number of those probes allows the fingerprinting of a target polynucleotide or the relative mapping of a collection of target polynucleotides, as described in greater detail below.

In the high resolution fingerprinting provided by a saturating collection of probes which include all possible subsequences of a given size, e.g., 10-mers, collating of all the subsequences and determination of specific overlaps will be derived and the entire sequence can usually be reconstructed.

Although a polynucleotide sequence analysis is a preferred embodiment, for which the specific reagents are most easily accessible, the invention is also applicable to analysis of other polymers, including polypeptides, carbohydrates, and synthetic polymers, including α -, β -, and ω -amino acids, polyurethanes, polyesters, polycarbonates, polyureas,

US 6,440,667 B1

7

polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, polyacetates, and mixed polymers. Various optical isomers, e.g., various D- and L- forms of the monomers, may be used.

Sequence analysis will take the form of complete sequence determination, to the level of the sequence of individual subunits along the entire length of the target sequence. Sequence analysis also takes the form of sequence homology, e.g., less than absolute subunit resolution, where "similarity" in the sequence will be detectable, or the form of selective sequences of homology interspersed at specific or irregular locations.

In either case, the sequence is determinable at selective resolution or at particular locations. Thus, the hybridization method will be useful as a means for identification, e.g., a "fingerprint", much like a Southern hybridization method is used. It is also useful to map particular target sequences.

B. VLSIPS™ Technology

The invention is enabled by the development of technology to prepare substrates on which specific reagents may be either positionally attached or synthesized. In particular, the very large scale immobilized polymer synthesis (VLSIPS™) technology allows for the very high density production of an enormous diversity of reagents mapped out in a known matrix pattern on a substrate. These reagents specifically recognize subsequences in a target polymer and bind thereto, producing a map of positionally defined regions of interaction. These map positions are convertible into actual features recognized, and thus would be present in the target molecule of interest.

As indicated, the sequence specific recognition reagents will often be oligonucleotides which hybridize with fidelity and discrimination to the target sequence. For use with other polymers, monoclonal or polyclonal antibodies having high sequence specificity will often be used.

In the generic sense, the VLSIPS technology allows the production of a substrate with a high density matrix of positionally mapped regions with specific recognition reagents attached at each distinct region. By use of protective groups which can be positionally removed, or added, the regions can be activated or deactivated for addition of particular reagents or compounds. Details of the protection are described below and in related Pirrung et al. (1992) U.S. Pat. No. 5,143,854. In a preferred embodiment, photosensitive protecting agents will be used and the regions of activation or deactivation may be controlled by electro-optical and optical methods, similar to many of the processes used in semiconductor wafer and chip fabrication.

In the nucleic acid nucleotide sequencing application, a VLSIPS substrate is synthesized having positionally defined oligonucleotide probes. See Pirrung et al. (1992) U.S. Pat. No. 5,143,854; and U.S. Ser. No. 07/624,120, now abandoned. By use of masking technology and photosensitive synthetic subunits, the VLSIPS apparatus allows for the stepwise synthesis of polymers according to a positionally defined matrix pattern. Each oligonucleotide probe will be synthesized at known and defined positional locations on the substrate. This forms a matrix pattern of known relationship between position and specificity of interaction. The VLSIPS technology allows the production of a very large number of different oligonucleotide probes to be simultaneously and automatically synthesized including numbers in excess of about 10^2 , 10^3 , 10^4 , 10^5 , 10^6 , or even more, and at densities of at least about 10^2 , $10^3/\text{cm}^2$, $10^4/\text{cm}^2$, $10^5/\text{cm}^2$ and up to $10^6/\text{cm}^2$ or more. This application discloses methods for synthesizing polymers on a silicon or other suitably derivatized substrate, methods and chemistry for synthesizing

8

specific types of biological polymers on those substrates, apparatus for scanning and detecting whether interaction has occurred at specific locations on the substrate, and various other technologies related to the use of a high density very large scale immobilized polymer substrate. In particular, sequencing, fingerprinting, and mapping applications are discussed herein in detail, though related technologies are described in simultaneously filed applications U.S. Ser. No. 07/624,120, now abandoned; and U.S. Ser. No. 07/517,659; Dower et al. (1995) U.S. Pat. No. 5,427,908, each of which is hereby incorporated herein by reference.

In other embodiments, antibody probes will be generated which specifically recognize particular subsequences found on a polymer. Antibodies would be generated which are specific for recognizing a three contiguous amino acid sequence, and monoclonal antibodies may be preferred. Optimally, these antibodies would not recognize any sequences other than the specific three amino acid stretch desired and the binding affinity should be insensitive to flanking or remote sequences found on a target molecule. Likewise, antibodies specific for particular carbohydrate linkages or sequences will be generated. A similar approach could be used for preparing specific reagents which recognize other polymer subunit sequences. These reagents would typically be site specifically localized to a substrate matrix pattern where the regions are closely packed.

These reagents could be individually attached at specific sites on the substrate in a matrix by an automated procedure where the regions are positionally targeted by some other specific mechanism, e.g., one which would allow the entire collection of reagents to be attached to the substrate in a single reaction. Each reagent could be separately attached to a specific oligonucleotide sequence by an automated procedure. This would produce a collection of reagents where, e.g., each monoclonal antibody would have a unique oligonucleotide sequence attached to it. By virtue of a VLSIPS substrate which has different complementary oligonucleotides synthesized on it, each monoclonal antibody would specifically be bound only at that site on the substrate where the complementary oligonucleotide has been synthesized. A crosslinking step would fix the reagent to the substrate. See, e.g., Dattagupta et al. (1985) U.S. Pat. No. 4,542,102 and (1987) U.S. Pat. No. 4,713,326; and Chatterjee, M. et al. (1990) *J. Am. Chem. Soc.* 112:6397-6399, which are hereby incorporated herein by reference. This allows a high density positionally specific collection of specific recognition reagents, e.g., monoclonal antibodies, to be immobilized to a solid substrate using an automated system.

The regions which define particular reagents will usually be generated by selective protecting groups which may be activated or deactivated. Typically the protecting group will be bound to a monomer subunit or spatial region, and can be spatially affected by an activator, such as electromagnetic radiation. Examples of protective groups with utility herein include nitroveratryl oxycarbonyl (NVOC), nitrobenzyl oxycarbonyl (NBOC), dimethyl dimethoxy benzyloxy carbonyl, 5-bromo-7-nitroindolyl, O-hydroxy- α -methyl cinnamoyl, and 2-oxymethylene anthraquinone. Examples of activators include ion beams, electric fields, magnetic fields, electron beams, x-ray, and other forms of electromagnetic radiation.

C. Binary Masking

In fact, the means for producing a substrate useful for these techniques are explained in Pirrung et al. (1992) U.S. Pat. No. 5,143,854, which is hereby incorporated herein by reference. However, there are various particular ways to optimize the synthetic processes. Many of these methods are described in Ser. No. 07/624,120, now abandoned.

US 6,440,667 B1

9

Briefly, the binary synthesis strategy refers to an ordered strategy for parallel synthesis of diverse polymer sequences by sequential addition of reagents which may be represented by a reactant matrix, and a switch matrix, the product of which is a product matrix. A reactant matrix is a 1xn matrix of the building blocks to be added. The switch matrix is all or a subset of the binary numbers from 1 to n arranged in columns. In preferred embodiments, a binary strategy is one in which at least two successive steps illuminate half of a region of interest on the substrate. In most preferred embodiments, binary synthesis refers to a synthesis strategy which also factors a previous addition step. For example, a strategy in which a switch matrix for a masking strategy halves regions that were previously illuminated, illuminating about half of the previously illuminated region and protecting the remaining half (while also protecting about half of previously protected regions and illuminating about half of previously protected regions). It will be recognized that binary rounds may be interspersed with non-binary rounds and that only a portion of a substrate may be subjected to a binary scheme, but will still be considered to be a binary masking scheme within the definition herein. A binary "masking" strategy is a binary synthesis which uses light to remove protective groups from materials for addition of other materials such as nucleotides or amino acids.

In particular, this procedure provides a simplified and highly efficient method for saturating all possible sequences of a defined length polymer. This masking strategy is also particularly useful in producing all possible oligonucleotide sequence probes of a given length.

D. Applications

The technology provided by the present invention has very broad applications. Although described specifically for polynucleotide sequences, similar sequencing, fingerprinting, mapping, and screening procedures can be applied to polypeptide, carbohydrate, or other polymers. In particular, the present invention may be used to completely sequence a given target sequence to subunit resolution. This may be for de novo sequencing, or may be used in conjunction with a second sequencing procedure to provide independent verification. See, e.g., (1988) *Science* 242:1245. For example, a large polynucleotide sequence defined by either the Maxam and Gilbert technique or by the Sanger technique may be verified by using the present invention.

In addition, by selection of appropriate probes, a polynucleotide sequence can be fingerprinted. Fingerprinting is a less detailed sequence analysis which usually involves the characterization of a sequence by a combination of defined features. Sequence fingerprinting is particularly useful because the repertoire of possible features which can be tested is virtually infinite. Moreover, the stringency of matching is also variable depending upon the application. A Southern Blot analysis may be characterized as a means of simple fingerprint analysis.

Fingerprinting analysis may be performed to the resolution of specific nucleotides, or may be used to determine homologies, most commonly for large segments. In particular, an array of oligonucleotide probes of virtually any workable size may be positionally localized on a matrix and used to probe a sequence for either absolute complementary matching, or homology to the desired level of stringency using selected hybridization conditions.

In addition, the present invention provides means for mapping analysis of a target sequence or sequences. Mapping will usually involve the sequential ordering of a plurality of various sequences, or may involve the localization of a particular sequence within a plurality of sequences. This

10

may be achieved by immobilizing particular large segments onto the matrix and probing with a shorter sequence to determine which of the large sequences contain that smaller sequence. Alternatively, relatively shorter probes of known or random sequence may be immobilized to the matrix and a map of various different target sequences may be determined from overlaps. Principles of such an approach are described in some detail by Evans et al. (1989) "Physical Mapping of Complex Genomes by Cosmid Multiplex Analysis," *Proc. Natl. Acad. Sci. USA* 86:5030-5034; Michiels et al. (1987) "Molecular Approaches to Genome Analysis: A Strategy for the Construction of Ordered Overlap Clone Libraries," *CABIOS* 3:203-210; Olsen et al. (1986) "Random-Clone Strategy for Genomic Restriction Mapping in Yeast," *Proc. Natl. Acad. Sci. USA* 83:7826-7830; Craig, et al. (1990) "Ordering of Cosmid Clones Covering the Herpes Simplex Virus Type I (HSV-I) Genome: A Test Case for Fingerprinting by Hybridization," *Nuc. Acids Res.* 18:2653-2660; and Coulson, et al. (1986) "Toward a Physical Map of the Genome of the Nematode *Caenorhabditis elegans*," *Proc. Natl. Acad. Sci. USA* 83:7821-7825; each of which is hereby incorporated herein by reference.

Fingerprinting analysis also provides a means of identification. In addition to its value in apprehension of criminals from whom a biological sample, e.g., blood, has been collected, fingerprinting can ensure personal identification for other reasons. For example, it may be useful for identification of bodies in tragedies such as fire, flood, and vehicle crashes. In other cases the identification may be useful in identification of persons suffering from amnesia, or of missing persons. Other forensics applications include establishing the identity of a person, e.g., military identification "dog tags", or may be used in identifying the source of particular biological samples. Fingerprinting technology is described, e.g., in Carrano, et al. (1989) "A High-Resolution, Fluorescence-Based, Semi-automated method for DNA Fingerprinting," *Genomics* 4: 129-136, which is hereby incorporated herein by reference. See, e.g., table I, for nucleic acid applications, and corresponding applications may be accomplished using polypeptides.

TABLE I

VLSIPS™ TECHNOLOGY IN NUCLEIC ACIDS	
I.	Construction of Chips
II.	Applications
A.	Sequencing
1.	Primary sequencing
2.	Secondary sequencing (sequence checking)
3.	Large scale mapping
4.	Fingerprinting
B.	Duplex/Triplex formation
1.	Antisense
2.	Sequence specific function modulation (e.g. promoter inhibition)
C.	Diagnosis
1.	Genetic markers
2.	Type markers
a.	Blood donors
b.	Tissue transplants
D.	Microbiology
1.	Clinical microbiology
2.	Food microbiology
III.	Instrumentation
A.	Chip machines
B.	Detection